

OPTIONAL QUESTIONNAIRE**SECTION C. BEVERAGES****Optional detailed information about alcohol drinking habits**

1. If interviewee reported to drink alcohol regularly at some point in life, answer separately for beer (1 can, 330ml), wine (1 glass, 125ml), and hard liquor (one pub measure, 25ml).

Beginning with when you started drinking alcohol regularly; please indicate any changes in alcohol consumption

1a. Period. FROM AGE _____ TO AGE _____ YEARS

1b. How many drinks did you usually drink per day/week/month? _____ DAY 1/

_____ WEEK 2 / _____ MONTH 3/ DK 8888

2. Aside from your usual pattern of drinking, did you ever regularly, but infrequently drink more than you usually do? YES 1 / NO 2 / DK 8888

3. At what age did you first experience periods of drinking more than you usually do?

_____ YEARS

4. At what age did you last experience periods of drinking more than you usually do?

_____ YEARS

5. On days that you drank more than you usually do, about how many drinks would you drink in one day? _____ DRINKS

6. About how many times per month or year would you drink a larger amount than you usually do? _____ MONTH 1/ _____ YEAR 2 / DK 8888

7. During how many months or years did you drink this larger amount? _____ MONTH 1/

_____ YEAR 2 / DK 8888

GUIDELINES FOR COMPLETING THE OPTIONAL SECTIONS OF THE QUESTIONNAIRE

This questionnaire was designed for case-control studies. Questionnaire must be applied by trained interviewers and questions must not be modified in any way.

Section C. Beverages

Optional detailed information about alcohol drinking habits

1. If the patient reports to ever drinking alcohol regularly, for every type of alcohol (beer/wine/hard liquor) indicate relevant changes in drinking habits
 - 1a. Indicate age of beginning and age of end for each period of specific consumption habits
 - 1b. Indicate the average number of drinks consumed by the patient either by day, week, or month during each period of specific consumption habits
2. Indicate if the patient ever had a period in which drank regularly but infrequently more than usual
3. Indicate the age at which the patient reported to start the periods of more than usual drinking
4. Indicate the age at which the patient reported to stop the periods of more than usual drinking
5. Indicate the average number of drinks consumed in one day (all types of alcohol) during the periods of more than usual drinking
6. Indicate the frequency (times a month/year) of the periods of more than usual drinking
7. Indicate the time (number of months/years) the periods of more than usual drinking lasted